

ヒト血清アルブミンの体内動態に及ぼす 表面電荷と α -ヘリックス含量の影響

岩尾康範*、平池美香子*、Ulrich Kragh-Hansen*、米良克美*、野口大志*、
安楽 誠、川井恵一*、丸山 徹*、小田切優樹*

Biochimica et Biophysica Acta, 1774(12), 1582-1590 (2007)

Changes of net charge and alpha-helical content affect the pharmacokinetic properties of human serum albumin

Yasunori Iwao*, Mikako Hiraike*, Ulrich Kragh-Hansen*, Katsumi Mera*,
Taishi Noguchi*, Makoto Anraku, Keiichi Kawai*, Toru Maruyama*, Masaki Otagiri*

ABSTRACT The pharmacokinetics of 17 genetic variants of human serum albumin with single-residue mutations and their corresponding normal albumin were studied in mice. In all cases, the plasma half-life was affected, but only variants with +2 changes in charge prolonged it, whereas changes in hydrophobicity decreased it. Good positive and negative correlations were found between changes in alpha-helical content taking place in domains I+III and domain II, respectively, and changes in half-lives. Liver and kidney uptake clearances were also modified: alpha-helical changes of domains I+III showed good negative correlations to both types of clearances, whereas changes in domain II only had a good positive correlation to kidney uptake clearance. The present information should be useful when designing recombinant, therapeutical albumins or albumin products with a modified plasma half-life.

抄録 ヒト血清アルブミンの17個の1残基変異体（バリエント）の体内動態を正常アルブミンと比較検討した。その結果、バリエントによる構造変化の中でもその表面電荷及び α -ヘリックス含量の変化と体内動態変化が相関することが明らかとなった。今回得られた結果はアルブミン製剤の設計において、重要な基礎資料として期待される。

* Graduate School of Medical and Pharmaceutical Sciences, Kumamoto University
熊本大学大学院医学薬学研究部